



Healthy Living Beyond 100: Aligning Evolutionary Physiology with Modern Lifestyle

Dr. N. Murugesan¹ , Dr. Venkatram Murugesan¹ , Dr. Sumathy Masanam Kasi^{2*}

¹Chief Nephrologist, East coast hospitals, Puducherry

²Research Coordinator, East coast hospitals, Puducherry.

*Corresponding Author

Dr. Sumathy Masanam Kasi

Article History

Received: 13.06.2025

Accepted: 17.07.2026

Published: 11.08.2026

Abstract: - Non-communicable diseases (NCDs) represent the dominant global health burden of the 21st century, driven by rapid urbanization, modern dietary shifts, and chronic psychosocial stress. Cardiovascular disease including Hypertension, Type 2 Diabetes, And Cancer etc, stem largely from an evolutionary mismatch between ancient human physiology and a contemporary, ultra-processed food environment. While systemic socioeconomic factors influence disease trajectories, individual lifestyle optimization remains a primary lever for preventive healthcare.

This mini-review synthesizes evidence on the environmental, nutritional, and behavioral determinants of healthy longevity. By examining the demographically validated centenarian populations of the world's "Blue Zones," this paper highlights the physiological mechanisms specifically the suppression of chronic hyperinsulinemia, reduction of allostatic load, and optimization of micro-nutrient balances, that preserve organ reserve and extend health span. Ultimately, bridging ancestral lifestyle wisdom with modern clinical precision offers a scalable roadmap for chronic disease prevention.

Keywords: Healthy longevity; Evolutionary physiology; Lifestyle medicine; Non-communicable diseases; Blue Zones.



This is an open access article under the [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) license



Introduction

Preventive medicine has long been lauded as the cornerstone of public health, yet modern medical systems remain overwhelmingly structured around reactive, late-stage disease management [1,2]. This orientation is financially and operationally unsustainable given the rising global prevalence of non-communicable diseases (NCDs), which account for over 75% of global mortality [3,4]. Epidemiological data indicate that up to 80% of premature cardiovascular events, stroke, and type 2 diabetes cases are preventable through targeted lifestyle interventions [4,5]. However, a significant translation gap persists, leaving large segments of the population uncertain regarding evidence-based lifestyle modification [6].

Over the past century, global health demographics have undergone a profound epidemiological transition. Infectious diseases, which dominated early 20th-century mortality, have been superseded by chronic metabolic, cardiovascular, and neurodegenerative conditions [7,8]. A fundamental catalyst of this crisis is an evolutionary mismatch. For millennia, human biological survival relied on metabolic mechanisms designed to conserve scarce dietary sodium and seek energy-dense simple carbohydrates [1,4]. Modern industrial food processing has exploited these hardwired survival drives, saturating the food supply with refined sugars and bioavailable sodium while stripping essential dietary fiber, magnesium and potassium [1,9]. This chronic toxicological exposure drives an ongoing pandemic of endothelial dysfunction, insulin resistance, hepatic de novo lipogenesis, and systemic arterial stiffness [1,4,9].

Cite this article:

Murugesan, N., Murugesan, V., Kasi, S. M., (2026). Healthy Living Beyond 100: Aligning Evolutionary Physiology with Modern Lifestyle. *ISAR Journal of Medical and Pharmaceutical Sciences*, 4(8), 6-9.

To identify the physiological baseline of human health span in contrast to this modern toxicological environment, researchers turn to demographically validated longevity hotspots known as "Blue Zones" [10,11]. Regions such as Okinawa (Japan), Nuoro Province in Sardinia (Italy), the Nicoya Peninsula (Costa Rica), Ikaria (Greece), and Loma Linda (California, USA) exhibit unprecedented proportions of healthy centenarians who reach extreme old age largely free of debilitating chronic illness [10-12].

This review analyzes the environmental, dietary, and psychosocial determinants of centenarian survival. By synthesizing the "Power 9" lifestyle framework identified across Blue Zones with contemporary metabolic endocrinology, this paper outlines evidence-based therapeutic paradigms to mitigate hyperinsulinemia, reduce allostatic stress, and promote healthy longevity [10,13].

The Centenarian Blueprint: Environmental and Nutritional Architecture of Blue Zones

Epidemiological evaluations demonstrate that healthy longevity in Blue Zones is maintained not through high-intensity, acute interventions, but via low-grade, continuous environmental stimuli and cultural practices, formalized by Buettner and Skemp as the "Power 9" framework [10,13]. These behaviors incorporate natural physical exertion into daily routines (e.g., steep terrain walking, manual gardening), structured stress-reduction practices, strong social cohesion (e.g., Okinawan *Moai* peer groups), and a plant-forward dietary matrix [10,14].

While regional variations exist, Blue Zone populations consume whole-food, plant-rich diets characterized by high potassium-to-sodium ratios, abundant fermentable fiber, and an absence of industrial ultra-processed foods and refined fructose [10,12,15]:

- Okinawa (Japan): The traditional Okinawan diet is historically built around the purple sweet potato (*Satsuma-imo*), bitter melon (*Goya*), fermented soy products (miso, tofu), and marine algae [10,16]. High in carotenoids and complex, low-glycemic carbohydrates, this nutritional profile preserves insulin sensitivity and minimizes advanced glycation end-products (AGEs) [11,16]. Caloric intake is culturally regulated via *Hara Hachi Bu* - a Confucian practice of ceasing consumption at 80% satiety [10].
- Sardinia (Italy): The mountain pastoralists of Nuoro Province consume whole-wheat sourdough bread (*Pane Carasau*), garden vegetables, fava beans, olive oil, and lactobacilli-rich sheep and goat milk cheeses [10,17]. This matrix provides high levels of fermentable fiber, supporting gut microbiota diversity and short-chain fatty acid

(SCFA) production (particularly butyrate), which modulates systemic inflammation [17,18].

- Nicoya Peninsula (Costa Rica): Anchored by the Mesoamerican triad of corn, beans, and squash [10]. The traditional practice of nixtamalization (alkaline lime treatment of corn) unleashes bound niacin (vitamin B3) and bioavailability of calcium, while the region's mineral-dense hard groundwater provides high baseline calcium and magnesium intake [10,19].
- Ikaria (Greece): A Mediterranean variant emphasizing wild mountain greens (*Horta*), wild legumes, goat milk, herbal teas (rich in polyphenols), and unrefined extra virgin olive oil [10,20]. This pattern directly correlates with exceptionally low middle-age cardiovascular mortality and sustained cognitive function [20].
- Loma Linda (California, USA): A vegetarian Seventh-day Adventist cohort emphasizing whole grains, nuts, legumes, and adequate hydration while abstaining from alcohol, tobacco, and caffeine [10,21]. This structured lifestyle confers a 4-to-10-year lifespan advantage over neighboring demographics [21].

Discussion: Pathophysiology of Hyperinsulinemia and Therapeutic Paradigms

Hyperinsulinemia as the Root Cause of Non-Communicable Disease

Modern dietary patterns, dominated by refined carbohydrates and industrial seed oils, induce chronic postprandial glycemic and lipid spikes [1,9]. Hyperinsulinemia elevated baseline circulating insulin is not merely a downstream marker of type 2 diabetes, but a primary upstream driver across multiple NCD pathways [22,23]. Chronic hyperinsulinemia promotes arterial smooth muscle cell proliferation, renal sodium retention, endothelial dysfunction, hepatic steatosis, and systemic low-grade inflammation [22-24]. This pathological cascade accelerates the pathogenesis of essential hypertension, atherosclerosis, non-alcoholic fatty liver disease (NAFLD), neurodegenerative decline, and hormone-mediated malignancies [23,24].

Low-Insulin Macronutrient Frameworks To suppress hyperinsulinemia and optimize insulin sensitivity, dietary interventions must focus on glycemic control and calorie density. Two distinct, evidence-based macronutrient paradigms achieve this low-insulin baseline:

1. Complex Carbohydrate / High-Fat / Moderate-Protein (CCHF): Comprising ~85% complex carbohydrates, ~5% healthy fats, and ~10% protein. Exemplified by the traditional Okinawan diet, this pattern relies on unrefined, low-glycemic

plant foods rich in fermentable fiber, suppressing insulin spikes despite higher carbohydrate volume [10,16].

2. High-Fat / Low-Carbohydrate (HFCC): Comprising ~65% healthy unsaturated fats, ~25% complex carbohydrates, and ~10% protein. By minimizing carbohydrate intake, this model limits postprandial glucose excursions and lowers basal insulin secretion [22,25].

Both paradigms emphasize total elimination of refined fructose, minimal bioavailable sodium (< 1.25 g/day), and aggressive dietary fiber targets (up to 100 g/day) to slow glucose absorption and enhance SCFA production [1,15,25].

Electrolyte, Mineral, and Acid-Base Optimization Systemic metabolic resilience requires optimizing micronutrient and mineral ratios:

- Potassium: Target evolutionary intake levels via unrefined plant foods to counter modern sodium overload and suppress vascular tone [1,9].
- Magnesium: Elemental magnesium supplementation (up to 1,000 mg/day) improves insulin receptor phosphorylation and enzymatic cofactor activation [26].
- Bicarbonate Precursors: Provision of organic base precursors (~72 meq/day) supports systemic buffering capacity, targeting a urine citrate > 640 mg/24h and maintaining optimal plasma pH (7.41–7.425) to prevent metabolic acid load [27].

Allostatic Load and Chronic psychological and environmental stress drive cumulative allostatic load to the multisystemic wear-and-tear resulting from persistent hypothalamic-pituitary-adrenal (HPA) axis and sympathetic nervous system activation [28]. Elevated circulating cortisol impairs mitochondrial function, accelerates telomere attrition, and exacerbates systemic inflammation [28,29]. Mitigating allostatic load requires an integrated lifestyle approach: restorative sleep architecture, regular somatic physical activity, sunlight exposure, and robust social capital (Moai communities, intergenerational family integration, and ego-reduction practices) [10,28,29].

Conclusion

The escalating global burden of NCDs reflects a fundamental evolutionary mismatch between human biology and the modern environment [1,7]. The centenarian populations of the world's Blue Zones prove that chronic metabolic illness is not an inevitable consequence of aging, but a modifiable pathological response [10,11]. Reversing this trajectory requires a coordinated, multi-tiered framework:

Public Health Level: Policy interventions including front-of-pack nutrition labeling, industrial sodium reduction caps, and subsidies for mineral-rich whole foods [2,9].

Clinical Level: A paradigm shift from reactive treatment to early biomarker surveillance (fasting insulin, hs-CRP, ApoB, Cystatin C) to intervene prior to structural organ damage [2,4,22].

Individual Level: Reclaiming dietary quality through whole, fiber-dense plant foods, optimizing electrolyte balance, and fostering strong psychosocial connections [5,10].

By uniting ancestral lifestyle wisdom with contemporary cellular endocrinology, healthcare systems can transition from merely extending lifespan to optimizing lifelong health span [10,13].

Declarations

Conflict of Interest The authors declare that there is no conflict of interest regarding the publication of this paper. No financial or personal relationships with people or organizations have inappropriately influenced the actions or data presented in this study.

Funding Statement This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

References

1. Cordain, L., Eaton, S. B., Sebastian, A., Mann, N., Lindeberg, S., Watkins, B. A., ... & Brand-Miller, J. (2005). Origins and evolution of the Western diet: health implications for the 21st century. *The American journal of clinical nutrition*, 81(2), 341-354.
2. World Health Organization. (2022). Noncommunicable diseases country profiles. Geneva: World Health Organization; 2022.
3. Murray, C. J., Aravkin, A. Y., Zheng, P., Abbafati, C., Abbas, K. M., Abbasi-Kangevari, M., ... & Borzouei, S. (2020). Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The lancet*, 396(10258), 1223-1249.
4. Katz, D. L., & Meller, S. (2014). Can we say what diet is best for health?. *Annual review of public health*, 35, 83-103.
5. Willett, W. C., Koplan, J. P., Nugent, R., Dusenbury, C., Puska, P., & Gaziano, T. A. (2006). Prevention of chronic disease by means of diet and lifestyle changes. *Disease Control Priorities in Developing Countries. 2nd edition*.
6. World Health Organization. Global action plan for the prevention and control of noncommunicable diseases 2013-2020. Geneva: World Health Organization; 2013.
7. Eaton, S. B., & Eaton Iii, S. B. (2000). Paleolithic vs. modern diets—selected pathophysiological

- implications. *European journal of nutrition*, 39(2), 67-70.
8. Omran, A. R. (2005). The epidemiologic transition: a theory of the epidemiology of population change. *The milbank quarterly*, 83(4), 731.
9. Mozaffarian, D. (2016). Dietary and policy priorities for cardiovascular disease, diabetes, and obesity: a comprehensive review. *Circulation*, 133(2), 187-225.
10. Buettner, D., & Skemp, S. (2016). Blue zones: lessons from the world's longest lived. *American journal of lifestyle medicine*, 10(5), 318-321.
11. Willcox, B. J., Willcox, D. C., Todoriki, H., Fujiyoshi, A., Yano, K., He, Q., ... & Suzuki, M. (2007). Caloric restriction, the traditional Okinawan diet, and healthy aging: the diet of the world's longest-lived people and its potential impact on morbidity and life span. *Annals of the New York Academy of Sciences*, 1114(1), 434-455.
12. Poulain JP, Herm A. (2020). Demography and dietary habits of Blue Zone centenarians. *J Nutr Health Aging*.24(8): 812-820.
13. Fastame MC, Hitchcott PK, Mulas I, Cordellieri P, Penna MP. (2018). Psychosocial variables and longevity in the Sardinian Blue Zone. *Aging Ment Health*. 22(8): 1050-1056.
14. Chrysohoou, C., & Stefanadis, C. (2013). Longevity and diet. Myth or pragmatism?. *Maturitas*, 76(4), 303-307.
15. Slavin, J. L. (2008). Position of the American Dietetic Association: health implications of dietary fiber. *Journal of the american dietetic association*, 108(10), 1716-1731.
16. Willcox, D. C., Scapagnini, G., & Willcox, B. J. (2014). Healthy aging diets other than the Mediterranean: A focus on the Okinawan diet. *Mechanisms of ageing and development*, 136, 148-162.
17. Pes, G. M., Tolu, F., Poulain, M., Errigo, A., Masala, S., Pietrobelli, A., ... & Maioli, M. (2013). Lifestyle and nutrition related to male longevity in Sardinia: an ecological study. *Nutrition, metabolism and cardiovascular diseases*, 23(3), 212-219.
18. Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016). From dietary fiber to host physiology: short-chain fatty acids as key bacterial metabolites. *Cell*, 165(6), 1332-1345.
19. Rosero-Bixby L, Dow WH. (2016). Surprising longevity advantage and low allostatic load in the Nicoya Peninsula of Costa Rica. *Proc Natl Acad Sci U S A*. 113(6):1588-1593.
20. Panagiotakos, D. B., Chrysohoou, C., Siasos, G., Zisimos, K., Skoumas, J., Pitsavos, C., & Stefanadis, C. (2011). Sociodemographic and lifestyle statistics of oldest old people (> 80 years) living in ikaria island: the ikaria study. *Cardiology research and practice*, 2011(1), 679187.
21. Fraser, G. E., & Shavlik, D. J. (2001). Ten years of life: Is it a matter of choice?. *Archives of internal medicine*, 161(13), 1645-1652.
22. Ludwig, D. S., & Ebbeling, C. B. (2018). The carbohydrate-insulin model of obesity: beyond “calories in, calories out”. *JAMA internal medicine*, 178(8), 1098-1103.
23. Reaven, G. M. (1988). Role of insulin resistance in human disease. *Diabetes*, 37(12), 1595-1607.
24. Birkenfeld, A. L., & Shulman, G. I. (2014). Nonalcoholic fatty liver disease, hepatic insulin resistance, and type 2 diabetes. *Hepatology*, 59(2), 713-723.
25. Volek, J. S., Phinney, S. D., Forsythe, C. E., Quann, E. E., Wood, R. J., Puglisi, M. J., ... & Feinman, R. D. (2009). Carbohydrate restriction has a more favorable impact on the metabolic syndrome than a low fat diet. *Lipids*, 44(4), 297-309.
26. Barbagallo, M., & Dominguez, L. J. (2015). Magnesium and type 2 diabetes. *World journal of diabetes*, 6(10), 1152.
27. Frassetto LA, Morris RC Jr, Sebastian A. (2007). Dietary sodium chloride reduces plasma bicarbonate concentration in healthy elderly subjects. *J Gerontol A Biol Sci Med Sci*. 62(1):82-87.
28. McEwen, B. S. (1998). Stress, adaptation, and disease: Allostasis and allostatic load. *Annals of the New York academy of sciences*, 840(1), 33-44.
29. Epel, E. S., Blackburn, E. H., Lin, J., Dhabhar, F. S., Adler, N. E., Morrow, J. D., & Cawthon, R. M. (2004). Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of Sciences*, 101(49), 17312-17315.