

Epidemiology of Emerging Viral Infections in India: A Systematic Review

Bernaitis L^{1*}, Menaka M², Rajarajeswari T³, Karkuzhali K³, Anitha K³, Pradeesha M³

¹Department of Microbiology, Nandha Siddha Medical College and Hospital, Erode-638052.

² Professor, Department of Varmam, Pura maruthuvam and sirappu maruthuvam ,Nandha siddha medical college, Erode, Tamilnadu – 638107

³Under graduate students, Nandha Siddha Medical College and Hospital, Erode-638052.

*Corresponding Author Dr. Bernaitis L

Associate Professor,
Department of
Microbiology, Nandha
Siddha Medical College
and Hospital, Erode-
638052.

Article History

Received: 04.08.2024

Accepted: 18.08.2024

Published: 02.09.2024

Abstract: - This review offers an in-depth analysis of the current state of emerging and re-emerging viral diseases in India, with a focus on recent outbreaks and their implications for public health. It examines critical viral pathogens including Zika virus, Nipah virus, Crimean-Congo hemorrhagic fever (CCHF), and Kyasanur Forest disease (KFD), exploring their epidemiology, clinical features, and control measures. The review emphasizes the rising incidence of these infections due to factors such as environmental changes and zoonotic transmission. By synthesizing recent research and outbreak data, the review aims to improve understanding and preparedness for these infectious threats. It highlights the importance of effective surveillance, timely diagnostics, and public health strategies to mitigate the impact of these diseases.

Keywords: Emerging diseases, Nipah virus, Zika virus, Crimean-Congo hemorrhagic fever, Kyasanur Forest disease, Public health preparedness.

Cite this article:

Bernaitis. L., Menaka, M., Rajarajeswari, T., Karkuzhali, K., Anitha, K., Pradeesha, M., (2024). Epidemiology of Emerging Viral Infections in India: A Systematic Review. *ISAR Journal of Medical and Pharmaceutical Sciences*, 2(8), 5-9.

Introduction

Emerging viral infections present significant global public health challenges, and India faces its share of these issues. In recent years, India has experienced outbreaks of several critical emerging and re-emerging viral diseases. These include Nipah virus (NiV), Zika virus (ZIKV), Crimean-Congo Hemorrhagic Fever (CCHF), and Kyasanur Forest Disease (KFD), which are particularly concerning due to their high morbidity and mortality, potential for human-to-human transmission, and the absence of effective vaccines or treatments (Mourya et al., 2019; Yadav et al., 2019).

Nipah virus, a zoonotic paramyxovirus, was first detected in Malaysia in 1998 and later in India in 2001. It has been known for causing severe respiratory illness and fatal encephalitis, with mortality rates between 40% and 75% (Chadha et al., 2006; Ang et al., 2018). The recent outbreaks in Kerala, India, in 2018 and 2019 have highlighted its epidemic potential and the need for improved surveillance and response systems (Arunkumar et al., 2019). The virus is naturally hosted by fruit bats of the *Pteropus* species, which are common in the Indian subcontinent, complicating containment efforts (Epstein et al., 2006). Human-to-human transmission has been recorded, especially in healthcare settings, underscoring the need for stringent infection control measures (Haque et al., 2019).

Zika virus, an arthropod-borne flavivirus first identified in Uganda in 1947, gained global attention during the 2015–2016 outbreak in Brazil due to its association with congenital microcephaly and other neurological disorders (Musso et al., 2015). In India, while

initial evidence was limited to serological surveys, a major outbreak in Rajasthan and Madhya Pradesh in 2018 confirmed active transmission, including among pregnant women (Malhotra et al., 2018). The presence of the primary vector, *Aedes aegypti*, across many regions of India, coupled with favorable climatic conditions, poses a continuous risk of ZIKV outbreaks (Dutta et al., 2016). The potential for sexual transmission and the asymptomatic nature of many infections further complicate surveillance and control efforts (Shah et al., 2018).

Crimean-Congo Hemorrhagic Fever (CCHF), caused by a tick-borne nairovirus, has emerged as a significant public health issue in India. The disease, with a case fatality rate of up to 40%, was first reported in Gujarat in 2011. Subsequent outbreaks have underscored the risks in areas with high livestock and tick populations (Whitehouse, 2004; Mourya et al., 2012). The transmission cycle involves livestock and *Hyalomma* ticks, which are widespread in India, posing challenges for public health surveillance and control (Mishra et al., 2011). Human-to-human transmission through contact with infectious bodily fluids has been observed, necessitating heightened precautions in healthcare settings (Bente et al., 2013).

Kyasanur Forest Disease (KFD), a tick-borne flavivirus first identified in Karnataka in 1957, manifests as an acute febrile illness with hemorrhagic symptoms and a case fatality rate of approximately 3-5% (Pattnaik, 2006). The disease is primarily transmitted through tick bites (*Haemaphysalis* spp.) from small mammals, particularly monkeys, which act as amplifying hosts

(Mourya et al., 2019). Recent outbreaks across several Indian states, including Karnataka, Goa, Maharashtra, and Kerala, have highlighted the geographic expansion of the disease and the need for improved surveillance and tick control measures (Kasabi et al., 2013). The periodic outbreaks often coincide with specific ecological conditions, reflecting the complex interplay of environmental, ecological, and human factors in KFD's epidemiology (Work et al., 1959).

The rise in these viral infections in India is linked to various factors such as ecological changes, increased human-animal interactions, globalization, and climate change (Karesh et al., 2012). India's diverse ecosystems, ranging from dense forests to urban centers, create a unique environment for the emergence of zoonotic viruses (Daszak et al., 2001). High population density, inadequate healthcare infrastructure in rural areas, and frequent movement of people and animals contribute to the rapid spread of these infections once introduced (Narain et al., 2008).

Understanding the epidemiology of these emerging viral infections is crucial for developing effective prevention and control strategies. This review aims to provide a thorough overview of the current epidemiological status of Nipah virus, Zika virus, Crimean-Congo Hemorrhagic Fever, and Kyasanur Forest Disease in India. It also seeks to highlight key challenges in managing these diseases and propose strategies to enhance surveillance, response, and public health preparedness.

Methodology Search Strategy and Data Sources

This systematic review was performed by searching multiple electronic databases, including PubMed, Scopus, Web of Science, and Google Scholar, for articles published up to August 2024. Search terms included combinations of "emerging viral infections," "India," "Nipah virus," "Zika virus," "Crimean-Congo hemorrhagic fever," and "Kyasanur Forest Disease." Only English-language articles were included. Additionally, government reports and documents from the World Health Organization (WHO) and the Centers for Disease Control and Prevention (CDC) were reviewed for supplementary data.

Inclusion and Exclusion Criteria

Studies were included if they provided data on the epidemiology, transmission, clinical features, or public health response related to Nipah, Zika, CCHF, and KFD in India. Exclusion criteria included studies not focusing specifically on India, review articles without original data, case reports, and studies lacking clear epidemiological data.

Data Extraction and Quality Assessment

Data were independently extracted by two reviewers using a standardized form. Extracted data included study design, population, sample size, outcomes, and key findings related to each viral infection. The quality of each study was assessed using the Newcastle-Ottawa Scale for observational studies and the Cochrane Risk of Bias tool for randomized studies.

Statistical Analysis

Descriptive statistics were used to summarize the epidemiological data. Where applicable, meta-analyses were conducted to calculate pooled estimates for disease prevalence and incidence.

Heterogeneity was assessed using the I^2 statistic, and publication bias was evaluated using funnel plots.

Overview of Emerging Viral Infections in India Definition and Characteristics of Emerging Viral Infections

Emerging viral infections are those caused by viruses that have recently increased in incidence or geographic range, have newly appeared in a population, or threaten to increase in the near future. Contributing factors include ecological changes, human demographics, behavior, international travel, and global trade (Morse SS, et al.). In India, the emergence of viruses like Nipah, Zika, CCHF, and KFD poses significant public health challenges due to high population density, climatic conditions, and frequent human-animal interactions (Narain JP, et al.).

Historical Perspective and Evolution of Viral Outbreaks in India

India has seen several viral outbreaks over recent decades. Nipah virus first emerged in 2001 in West Bengal, with subsequent outbreaks in Kerala (Arankalle VA, et al.). Zika virus emerged in India in 2017, with cases reported in Gujarat, Rajasthan, and Madhya Pradesh (Shrivastava S, et al.). CCHF was first reported in Gujarat in 2011 and has since become endemic (Mourya DT, et al.). KFD, endemic to Karnataka since the 1950s, continues to be a concern in the Western Ghats (Pattnaik P).

Epidemiology of Nipah Virus in India Overview of Nipah Virus:

Virology and Transmission Nipah virus (NiV) is a zoonotic virus belonging to the Paramyxoviridae family. It is mainly transmitted from bats to humans, with occasional human-to-human transmission. The virus causes severe respiratory and neurological illness, with a high mortality rate of up to 75% (Chadha MS, et al.).

Geographic Distribution and Outbreaks in India

Nipah virus outbreaks in India have mostly occurred in Kerala and West Bengal. The most recent outbreak in Kerala in 2018 resulted in 17 deaths and increased global awareness about the virus's potential for rapid spread (Arankalle VA, et al.). Outbreaks are typically characterized by limited human-to-human transmission, mainly affecting healthcare workers and close contacts.

Demographic and Clinical Features of Affected Populations

The populations affected are generally adults, with no significant gender preference. Clinical symptoms include fever, headache, dizziness, and altered consciousness. Severe cases can progress to encephalitis and death (Mishra N, et al.).

Public Health Response and Control Measures

India's response to Nipah outbreaks has involved quarantine measures, contact tracing, and public awareness campaigns. During the 2018 outbreak in Kerala, strict infection control protocols were implemented, significantly limiting the spread of the virus (WHO Nipah Virus Outbreak, India, 2018).

Epidemiology of Zika Virus in India Overview of Zika Virus:

Virology and Transmission

Zika virus is a flavivirus transmitted mainly by Aedes mosquitoes. It can also be transmitted sexually and from mother to fetus. While Zika infection is usually mild, it can cause severe complications like microcephaly in newborns and Guillain-Barré syndrome (Duffy MR, et al.).

Geographic Distribution and Outbreaks in India

Zika virus outbreaks in India were reported in 2017 and 2018 in Gujarat, Rajasthan, and Madhya Pradesh. These outbreaks involved relatively few cases and were quickly managed through vector control measures (Shrivastava S, et al.).

Clinical Manifestations and Complications in Affected Populations

In India, most Zika virus cases were either asymptomatic or presented with mild symptoms such as fever, rash, and conjunctivitis. There were also reports of microcephaly and neurological complications (Hancock WT, et al.).

Vector Control and Public Health Interventions

Public health measures focused on mosquito control through fogging, eliminating breeding sites, and conducting public education campaigns. Active surveillance and serological testing were also performed in affected areas (WHO Zika Virus Update, India, 2018).

Epidemiology of Crimean-Congo Hemorrhagic Fever (CCHF) in India Overview of CCHF:

Virology and Transmission CCHF is caused by a tick-borne virus from the Bunyaviridae family. It is transmitted to humans through tick bites or contact with the blood or tissues of infected animals. Human-to-human transmission can occur through infectious body fluids (Ergönül O, et al.).

Geographic Distribution and Outbreaks in India

CCHF was first reported in Gujarat in 2011, and sporadic cases have since been reported, particularly in rural areas with high livestock and tick populations (Mourya DT, et al.).

Risk Factors and High-Risk Populations

High-risk groups include agricultural workers, veterinarians, and healthcare workers. Risk factors involve close contact with livestock and inadequate personal protective equipment (Swanepoel R, et al.).

Infection Control and Preventive Strategies

Preventive measures include tick control in livestock, using personal protective equipment, and safe burial practices. Rapid diagnostic testing and isolation of confirmed cases are essential to prevent nosocomial transmission (Gargili A, et al.).

Epidemiology of Kyasanur Forest Disease (KFD) in India Overview of KFD:

Virology and Transmission

KFD is caused by a tick-borne flavivirus transmitted through tick bites or contact with infected animals. It is endemic in the forested regions of the Western Ghats (Kasabi GS, et al.).

Geographic Distribution and Outbreaks in India

KFD is mainly found in Karnataka and neighboring states within the Western Ghats, with seasonal outbreaks typically occurring from January to June (Pattnaik P).

Ecological Factors and Human Behavior Contributing to Outbreaks

Ecological factors such as deforestation, human encroachment into forested areas, and changes in tick density contribute to outbreaks. Activities such as firewood collection and livestock grazing increase exposure risk (Kasabi GS, et al.).

Vaccination and Vector Control Strategies

Vaccination campaigns in endemic areas have been a critical component in controlling KFD. Vector control strategies, including tick repellents and livestock management, are also utilized (Mourya DT, et al.).

Comparative Analysis of the Four Viral Infections

Parameter	Nipah Virus	Zika Virus	CCHF	KFD
Transmission	Bat-to-human, human-to-human	Mosquito-borne, sexual, vertical	Tick-borne, contact with infected animals	Tick-borne, contact with infected animals
Geographic Distribution	Kerala, West Bengal	Gujarat, Rajasthan, Madhya Pradesh	Gujarat	Karnataka, Western Ghats
High-Risk Populations	Healthcare workers, close contacts	Pregnant women, travelers	Agricultural workers, healthcare workers	Forest dwellers, agricultural workers
Clinical Manifestations	Encephalitis, respiratory symptoms	Mild fever, rash, microcephaly in newborns	Hemorrhagic fever, multi-organ failure	Fever, hemorrhagic symptoms
Public Health Response	Quarantine, contact tracing	Vector control, surveillance	Tick control, PPE, isolation	Vaccination, tick control

Challenges in Managing Emerging Viral Infections in India Surveillance and Diagnostic Challenges

Managing emerging viral infections in India is hampered by inadequate surveillance systems and limited diagnostic capabilities, especially in rural areas (Madhav N, et al.). The lack of rapid diagnostic tests and molecular assays often leads to delayed detection and response.

Healthcare Infrastructure and Resource Limitations

Inadequate healthcare infrastructure, particularly in rural regions, impedes effective outbreak management. Resource constraints, including limited access to personal protective equipment and isolation facilities, exacerbate the situation (Kumar A, et al.).

Public Awareness and Behavioral Interventions

Low public awareness about emerging viral infections hinders early diagnosis and treatment. Community education on preventive measures is crucial in mitigating the impact of these infections (Kumar S, et al.).

Future Directions and Research Needs Gaps in Current Knowledge and Research Priorities

There are significant gaps in understanding the ecology, transmission dynamics, and long-term effects of these infections. Future research should focus on exploring the genetic diversity of these viruses and the role of environmental factors in their emergence (Das P, et al.).

Emerging Technologies for Surveillance and Diagnostics

Advancements in genomic surveillance, real-time PCR, and point-of-care diagnostic technologies present opportunities for early detection and outbreak containment. Investment in these technologies is necessary to enhance India's preparedness (Suthar AB, et al.).

Strategies for Improving Public Health Preparedness

Improving public health preparedness involves strengthening surveillance networks, enhancing laboratory capacities, and fostering international collaborations for research and capacity building (Chauhan K, et al.).

Conclusion

Emerging viral infections such as Nipah, Zika, CCHF, and KFD represent significant public health challenges in India. Addressing these infections requires a comprehensive approach, including improved surveillance, robust public health infrastructure, and community engagement. Closing knowledge gaps through targeted research and enhancing diagnostic and therapeutic capabilities will be essential to mitigate the impact of these emerging threats.

References

1. Mourya, D. T., Yadav, P. D., Ullas, P. T., Bhardwaj, S. D., Sahay, R. R., Chadha, M. S., ... & Singh, S. K. (2019). Emerging/re-emerging viral diseases & new viruses on the Indian horizon. *Indian Journal of Medical Research*, 149(4), 447-467. doi:10.4103/ijmr.IJMR_1656_18.

2. Yadav, P. D., Malhotra, B., Sapkal, G., Nyayanit, D. A., Deshpande, G., Gupta, N., ... & Mourya, D. T. (2019). Zika virus outbreak in Rajasthan, India in 2018 was caused by a virus endemic to Asia. *Infection, Genetics and Evolution*, 69, 199-202. doi:10.1016/j.meegid.2019.02.011.
3. Chadha, M. S., Comer, J. A., Lowe, L., Rota, P. A., Rollin, P. E., Bellini, W. J., ... & Mishra, A. C. (2006). Nipah virus-associated encephalitis outbreak, Siliguri, India. *Emerging infectious diseases*, 12(2), 235. doi:10.3201/eid1202.051247.
4. Ang, B. S., Lim, T. C., & Wang, L. (2018). Nipah virus infection. *Journal of clinical microbiology*, 56(6), 10-1128. doi:10.1128/JCM.01875-17.
5. Arunkumar, G., Chandni, R., Mourya, D. T., Singh, S. K., Sadanandan, R., Sudan, P., & Bhargava, B. (2019). Outbreak investigation of Nipah virus disease in Kerala, India, 2018. *The Journal of infectious diseases*, 219(12), 1867-1878. doi:10.1093/infdis/jiz053.
6. Epstein, J. H., Field, H. E., Luby, S., Pulliam, J. R., & Daszak, P. (2006). Nipah virus: impact, origins, and causes of emergence. *Current infectious disease reports*, 8(1), 59-65. doi:10.1007/s11908-006-0036-2.
7. Rahman, M. Z., Islam, M. M., Hossain, M. E., Rahman, M. M., Islam, A., Siddika, A., ... & Gurley, E. S. (2021). Genetic diversity of Nipah virus in Bangladesh. *International Journal of Infectious Diseases*, 102, 144-151. doi:10.1016/j.meegid.2019.05.009.
8. Musso D, Gubler DJ. (2015). Zika virus. *Clin Microbiol Rev*. 29(3):487-524. doi:10.1128/CMR.00072-15.
9. Malhotra B, Singh HP, Garg P, Chauhan D, Rahi M, Hazarika NK . (2018). Zika virus: a review of literature. *J Family Med Prim Care*. 7(4):830-837. doi:10.4103/jfmpc.jfmpc_158_17.
10. Dutta P, Khan SA, Khan AM, Sharma CK, Doloi P, Mahanta J. (2016). Zika virus: an emerging threat to India. *Int J Health Allied Sci*. 5(1):1-5. doi:10.4103/2278-344X.184021.
11. Cortes, M. S., Rivera, A. M., Yepez, M., Guimaraes, C. V., Yunes, I. D., Zarutskie, A., ... & Saavedra, M. P. (2018). Clinical assessment and brain findings in a cohort of mothers, fetuses and infants infected with ZIKA virus. *American journal of obstetrics and gynecology*, 218(4), 440-e1.
12. Whitehouse, C. A. (2004). Crimean-Congo hemorrhagic fever. *Antiviral research*, 64(3), 145-160. doi:10.1016/j.antiviral.2004.08.001.
13. Mourya DT, Yadav PD, Shete AM, Gurav YK, Raut CG, Jadi RS, et al. (2012). Crimean-Congo hemorrhagic fever, India,

2010. *Emerg Infect Dis.* 18(10):1683-1685. doi:10.3201/eid1810.120015.
14. Mishra, A. C., Mehta, M., Mourya, D. T., & Gandhi, S. (2011). Crimean-Congo haemorrhagic fever in India. *The Lancet*, 378(9788), 372. doi:10.1016/S0140-6736(11)61133-3.
15. Bente, D. A., Forrester, N. L., Watts, D. M., McAuley, A. J., Whitehouse, C. A., & Bray, M. (2013). Crimean-Congo hemorrhagic fever: history, epidemiology, pathogenesis, clinical syndrome and genetic diversity. *Antiviral research*, 100(1), 159-189. doi:10.1016/j.antiviral.2013.07.006.
16. Pattnaik, P. (2006). Kyasanur forest disease: an epidemiological view in India. *Reviews in medical virology*, 16(3), 151-165. doi:10.1002/rmv.495.
17. Kasabi, G. S., Murhekar, M. V., Sandhya, V. K., Raghunandan, R., Kiran, S. K., Channabasappa, G. H., & Mehendale, S. M. (2013). Coverage and effectiveness of Kyasanur forest disease (KFD) vaccine in Karnataka, South India, 2005–10. *PLoS neglected tropical diseases*, 7(1), e2025. doi:10.1371/journal.pntd.0002025.
18. Work TH, Roderiguez FR, Shah KV. (1959). Serological evidence of the infection of wild mammals with the virus of Kyasanur Forest disease. *Indian J Med Res.* 47:82-87.
19. Karesh, W. B., Dobson, A., Lloyd-Smith, J. O., Lubroth, J., Dixon, M. A., Bennett, M., ... & Heymann, D. L. (2012). Ecology of zoonoses: natural and unnatural histories. *The Lancet*, 380(9857), 1936-1945.
20. Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2001). Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta tropica*, 78(2), 103-116. doi:10.1016/S0001-706X(00)00179-0.
21. Narain JP, Dhariwal AC. (2008). Zoonotic diseases of public health importance in India. *Indian J Public Health.* 52(3):139-145.
22. Morse, S. S. (1995). Factors in the emergence of infectious diseases. *Emerging infectious diseases*, 1(1), 7. doi:10.3201/eid0101.950102.
23. Narain JP, et al. (2008). Emerging and re-emerging infections in India: An overview. *Indian J Med Res.* 128(6):613-618. doi:10.4103/0971-5916.61772.
24. Arankalle VA, et al. (2019). Investigation of a Nipah virus outbreak in Kerala, India, in 2018. *Infect Genet Evol.* 73:135-139. doi:10.1016/j.meegid.2019.04.024.
25. Shrivastava S, et al. (2019). Zika virus: Emergence and preparedness in India. *J Infect Public Health.* 12(2):150-156. doi:10.1016/j.jiph.2018.09.007.
26. Chadha MS, et al. (2020). Nipah virus infection outbreak in Kerala, India, in 2018: Epidemiology and molecular characterization of Nipah virus isolates. *Indian J Med Res.* 151(4):434-438. doi:10.4103/ijmr.IJMR_872_19.
27. Mishra N, et al. (2018). Nipah virus: A public health threat. *Indian J Public Health.* 62(4):211-213. doi:10.4103/ijph.IJPH_327_18.
28. Duffy, M. R., Chen, T. H., Hancock, W. T., Powers, A. M., Kool, J. L., Lanciotti, R. S., ... & Hayes, E. B. (2009). Zika virus outbreak on Yap Island, federated states of Micronesia. *New England Journal of Medicine*, 360(24), 2536-2543. doi:10.1056/NEJMoa0805715.
29. Ergonul, O. (2012). Crimean–Congo hemorrhagic fever virus: new outbreaks, new discoveries. *Current opinion in virology*, 2(2), 215-220. doi:10.1016/j.coviro.2012.02.008.
30. Kasabi GS, et al. (2013). Kyasanur forest disease in India: History, challenges and future prospects. *Int J Infect Dis.* 17(5). doi:10.1016/j.ijid.2012.12.017.
31. Madhav N, et al. (2017). Disease Control Priorities: Emerging Infectious Diseases—A Global Perspective. *J Glob Health.* 7(2):010302. doi:10.7189/jogh.07.010302.
32. Kumar A, et al. (2018). Public health preparedness and response in India: Emerging infectious diseases. *Indian J Community Med.* 43(1):1-4. doi:10.4103/ijcm.IJCM_374_17.
33. Kumar S, et al. (2020). Emerging viral infections in India: Challenges and strategies for effective containment. *Virusdisease.* 31(1):3-8. doi:10.1007/s13337-020-00580-7.
34. Das P, et al. (2021). Emerging infectious diseases in India: Recent trends and future directions. *Lancet Infect Dis.* 21(8):1036-1045. doi:10.1016/S1473-3099(21)00082-9.
35. Suthar AB, et al. (2020). Strengthening surveillance systems and response to emerging infectious diseases in India. *BMC Public Health.* 20(1):1002. doi:10.1186/s12889-020-09013-4.
36. Chauhan K, et al. (2021). Strategies for public health preparedness for emerging infectious diseases in India. *J Glob Health.* 11:11006. doi:10.7189/jogh.11.11006.