

Mindful Drawing: A Creative Approach to Alleviating Depressive Symptoms

Tianchan Chen¹, Yinan Zhang^{2*} 

¹Postgraduate, Massachusetts Institute of Technology, Room 905, Unit 6, Building 1, Longhu Tangning one, No.16 Zhongguancun East Road, Haidian District, Beijing, China

²Postgraduate, Peking University, Room 108, Philosophy Building, Peking University, No.5 Summer Palace Road, Haidian District, Beijing, China.

***Corresponding Author**
Yinan Zhang

Postgraduate, Peking
University.

Email: yinan@stu.pku.edu.cn

Article History

Received: 18.09.2024

Accepted: 14.10.2024

Published: 23.11.2024

Abstract: This study explores the effectiveness of mindful drawing as an intervention for alleviating depressive symptoms. Depression is a widespread mental health issue that significantly impacts individuals' quality of life. Current treatment methods, including pharmacotherapy and psychotherapy, have limitations in terms of side effects and accessibility. Mindful drawing, which combines mindfulness practice with artistic creation, may offer a promising non-invasive, low-cost alternative. This randomized controlled trial aims to evaluate the impact of mindful drawing on depressive symptoms, emotional regulation, and life satisfaction. Results are anticipated to show significant reductions in depressive symptoms and improvements in emotional regulation and overall mental health in the intervention group compared to the control group. These findings could provide valuable evidence supporting the use of mindful drawing as an adjunctive treatment for depression, especially for individuals resistant to traditional therapies.

Keywords: Mindful Drawing, Depression, Emotional Regulation, Art Therapy, Mental Health.

Cite this article:

Chen, T., Zhang, Y., (2024). Mindful Drawing: A Creative Approach to Alleviating Depressive Symptoms. *ISAR Journal of Arts, Humanities and Social Sciences*, 2(11), 111-115.

1. Introduction

1.1 Research Background

Depression is a prevalent global mental health issue that significantly impacts individuals' quality of life, social functioning, and work efficiency (WHO, 2017). According to the World Health Organization, depression is one of the leading causes of disability and reduced quality of life, and in severe cases, it may even lead to suicide (WHO, 2017). Although current interventions primarily involve pharmacological treatment and cognitive-behavioral therapy, these approaches have certain limitations in terms of long-term effectiveness and side effects (Cuijpers et al., 2016). In recent years, mindfulness interventions, as a form of non-pharmacological treatment, have garnered increasing attention and have been demonstrated to have significant effects in alleviating depressive symptoms (Kabat-Zinn, 1990; Segal et al., 2002).

Mindfulness is a practice that involves intentionally focusing one's attention on the present moment and observing internal and external experiences non-judgmentally, thereby regulating emotional responses, reducing rumination on negative emotions, and ultimately improving mental health (Steger et al., 2006). Mindful drawing is an intervention that combines mindfulness practice with artistic creation, guiding individuals to focus on the creative process, thereby enhancing self-expression,

reducing negative emotional reactions, and helping patients regulate their emotions (King, 2015). This approach not only carries the advantages of mindfulness interventions but also offers a non-verbal mode of expression, enabling patients to better understand and express their emotions, which is particularly useful for those who struggle to articulate their feelings verbally (Gussak, 2007).

Although mindful drawing has been increasingly applied in mental health interventions, empirical research on its efficacy remains limited, particularly regarding its impact on depression. Therefore, this study aims to explore the effectiveness of mindful drawing in alleviating depressive symptoms, providing new evidence and methodological support for adjunctive treatment of depression.

1.2 Research Significance

The significance of this study lies in exploring the effectiveness of mindful drawing in alleviating depressive symptoms and validating its efficacy through empirical data, thereby providing a new auxiliary treatment for individuals with depression. While current treatment methods, such as medication and psychotherapy, are effective, they often involve significant side effects or are difficult to access (Cuijpers et al., 2016). As a non-invasive and low-cost intervention, mindful drawing can help

alleviate depressive symptoms by enhancing emotional regulation, reducing negative rumination, and improving self-expression. This approach can complement traditional treatment, especially for those unwilling or unable to undergo pharmacological treatment, expanding the scope of depression intervention and advancing mental health intervention practices. Additionally, this study can provide a theoretical basis and practical guidance for clinical psychology and art therapy, facilitating the application of mindfulness and art therapy to promote comprehensive mental health improvement.

1.3 Research Objectives and Hypotheses

The main objective of this study is to investigate whether mindful drawing, as a psychological intervention, can effectively reduce depressive symptoms and improve overall mental health. Specifically, the study aims to evaluate the effectiveness of mindful drawing by comparing changes in depressive symptoms between the intervention and control groups before and after the intervention. This study hypothesizes that mindful drawing can significantly reduce depressive symptoms, enhance emotional regulation, and subsequently improve mental health. The specific hypotheses are:

Hypothesis 1 (H1): Depressive symptom scores in the intervention group will be significantly lower post-intervention compared to pre-intervention.

Hypothesis 2 (H2): The degree of reduction in depressive symptoms post-intervention will be greater in the mindful drawing group compared to the control group.

Hypothesis 3 (H3): Mindful drawing intervention will significantly improve participants' emotional regulation abilities and life satisfaction.

By testing these hypotheses, the study aims to further understand the mechanism of mindful drawing in depression intervention and provide theoretical support and empirical evidence for its clinical application.

2. Literature Review

2.1 The Current State of Depression and Treatment Approaches

Depression is a common and complex mental disorder that affects hundreds of millions of people globally (WHO, 2017). It not only affects individuals' emotional states but also negatively impacts social functioning, job performance, and interpersonal relationships (Ferrari et al., 2013). Current primary treatments for depression include pharmacotherapy and psychotherapy. Pharmacotherapy often involves antidepressants, such as selective serotonin reuptake inhibitors (SSRIs), which alleviate depressive symptoms by modulating neurotransmitters (Keller et al., 2000). However, pharmacotherapy can come with side effects such as weight gain, fatigue, and drug dependency, and its efficacy varies among individuals, with some patients showing poor response to medication (Jakobsen et al., 2017).

In terms of psychotherapy, cognitive-behavioral therapy (CBT) and mindfulness-based cognitive therapy (MBCT) are the most commonly used methods, particularly demonstrating effectiveness in the intervention of moderate to severe depression (Beck, 2011; Segal et al., 2002). CBT helps patients alleviate depressive symptoms by identifying and changing negative thought

patterns, while MBCT combines mindfulness practice to reduce reactions to negative emotions and thoughts (Segal et al., 2002). Despite their proven efficacy, psychotherapy is often less accessible for some patients, and it may take a considerable amount of time before noticeable results are achieved (Cuijpers et al., 2016).

Moreover, existing treatment approaches also exhibit shortcomings in terms of long-term efficacy and relapse prevention. Many patients experience a recurrence of symptoms after treatment discontinuation, making the search for more effective, long-term, and sustainable treatment methods a continuing challenge in clinical practice (Paykel, 2009). Therefore, combining innovative therapeutic approaches, such as art therapy and mindfulness interventions, may offer new directions and more comprehensive solutions for depression treatment (King, 2015).

2.2 The Theoretical Basis of Mindfulness Intervention

Mindfulness is a psychological practice originating from Eastern meditation traditions that involves intentionally focusing one's attention on the present moment and non-judgmentally observing internal and external experiences. Mindfulness practice can help individuals increase awareness of their emotions and thoughts, thereby better coping with stress and negative emotions (Kabat-Zinn, 1990). In mental health, mindfulness has been widely applied in the intervention of depression, anxiety, chronic pain, and various psychological and physical health issues (Baer, 2003).

The core mechanism of mindfulness intervention lies in alleviating the impact of negative emotions by improving emotional regulation (Gross, 2015). Emotional regulation refers to the processes by which individuals initiate, maintain, modulate, and alter their emotions. Mindfulness intervention reduces overreactivity to negative emotions, increases tolerance to negative feelings, and thereby decreases rumination and anxiety (Chambers et al., 2009). Moreover, mindfulness helps individuals observe and accept their emotional experiences, reducing judgment of negative emotions and improving overall mental health (Keng et al., 2011).

Recent years have seen promising results for mindfulness interventions in treating depression, particularly in mindfulness-based cognitive therapy (MBCT), which significantly reduces the risk of relapse by combining mindfulness practices with cognitive-behavioral techniques (Segal et al., 2002). Studies have demonstrated that mindfulness practices can effectively reduce negative emotional experiences in individuals with depression, improving their quality of life and emotional regulation abilities (Steger et al., 2006). Thus, mindfulness interventions, particularly in their influence on emotional regulation, are an area of significant interest in the field of psychology.

2.3 Art Therapy and Mindful Drawing

Art therapy is a therapeutic approach that promotes emotional expression and psychological healing through artistic creation. Its history can be traced back to the mid-20th century, when it was recognized as an effective method for helping individuals express emotions and resolve internal conflicts, particularly for those who find it difficult to articulate emotions verbally (King, 2015). The core concept of art therapy is that the creative process allows individuals to express themselves non-verbally, which is particularly important for individuals with depression, who often experience difficulties in emotional expression and suppression (Gussak, 2007).

Studies have shown that art therapy has a positive effect in reducing depressive symptoms. For example, Gussak (2007) found that prisoners participating in art therapy showed significant reductions in depressive symptoms. The creative process enables patients to externalize emotions, making it easier to identify and manage negative feelings, which is critical in alleviating depressive symptoms (King, 2015).

Mindful drawing, as an intervention that combines mindfulness practice with artistic creation, utilizes the strengths of both approaches by incorporating mindfulness into the creative process. Specifically, mindful drawing encourages individuals to focus on the act of drawing, paying attention to the movement of lines, colors, and shapes, which is similar to mindfulness practices involving breath focus or body scanning (Cohen & Sherman, 2014). Through this process, mindful drawing not only helps reduce rumination on negative emotions but also enhances self-efficacy and emotional regulation through creative expression (Monti et al., 2006).

Additionally, mindful drawing involves facilitating the experience of "flow" during the creative process—a state of complete immersion in an activity, which has been found to have positive effects on mental health (Csikszentmihalyi, 1990). By promoting emotional release and psychological balance through mindfulness during artistic creation, mindful drawing holds significant value for use as an adjunctive intervention in depression treatment (King, 2015).

3. Methodology

3.1 Research Design

This study employs a randomized controlled trial (RCT) design, with participants randomly assigned to either an intervention group or a control group. The intervention group will receive mindful drawing sessions, while the control group will receive routine psychological treatment. Participants in the intervention group will engage in mindful drawing activities twice a week, each lasting 90 minutes, for a total duration of eight weeks. The control group will receive standard psychological treatments, such as cognitive-behavioral therapy or supportive therapy (Kazdin, 2011).

The advantage of the RCT design lies in the ability to minimize systematic bias between intervention and control groups through randomization, ensuring a more scientific and valid comparison (Campbell & Stanley, 2015). By setting a control group, the unique effect of mindful drawing can be effectively evaluated, ensuring the internal validity of the experiment and the rationality of causal inference (Kazdin, 2011).

During the intervention period, all participants will be assessed for depressive symptoms at three time points: before the intervention (T1), immediately after the intervention (T2), and at follow-up (T3). Assessments will be conducted using the Patient Health Questionnaire (PHQ-9) and other measures, such as an emotional regulation scale (Kroenke et al., 2001). Data collected at multiple time points will help evaluate both the immediate and long-term effects of mindful drawing intervention on depressive symptoms.

3.2 Participants

Participants in this study will include individuals diagnosed with depression, aged 18-65 years. Diagnosis will be based on the

criteria of the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5). Exclusion criteria include individuals with severe mental illnesses, such as schizophrenia or bipolar disorder, and those unable to participate in the study. All participants will sign informed consent forms (World Medical Association, 2013).

3.3 Intervention

Participants in the intervention group will receive mindful drawing sessions twice a week for eight weeks, with each session lasting 90 minutes. Mindful drawing combines mindfulness practice and drawing creation, aimed at helping participants focus on their present experience during the drawing process, thereby reducing negative emotional responses (Peterson & Rappaport, 2014). Each session will have a specific theme and set of instructions:

Weeks 1-2: Exploration and Self-Connection

Instructions: "Freely draw your current feelings on paper. Do not worry about quality—just focus on each movement and your emotions."

Goal: Help participants begin to recognize their emotional state and express it through drawing.

Weeks 3-4: Colors of Emotion

Instructions: "Choose a color that represents your current emotion and express it through the use of color and shape."

Goal: Enhance participants' ability to recognize and accept emotions through the use of colors.

Weeks 5-6: Symbols of Inner Strength

Instructions: "Draw a symbol of your inner strength—it can be abstract or representational. Focus on your inner power and hope."

Goal: Help participants discover and depict their inner strength to increase confidence in coping with depression.

Weeks 7-8: Vision of the Future

Instructions: "Draw your hopes or vision for the future. This can take any form, as long as it represents your aspirations."

Goal: Encourage participants to express positive expectations for the future, fostering hope.

During the drawing sessions, therapists will provide a quiet and safe environment and offer non-directive support and encouragement as needed. Participants in the control group will receive routine psychological treatment, including psychological counseling and CBT. Both interventions will be conducted by professionally trained therapists.

3.4 Data Collection

Depressive symptoms will be assessed using the Patient Health Questionnaire-9 (PHQ-9), which is widely used to assess depressive symptoms and has demonstrated good reliability and validity (Kroenke et al., 2001). Data collection will occur at three time points: pre-intervention (T1), post-intervention (T2), and four weeks post-intervention follow-up (T3). Participants will complete the questionnaire under the guidance of a research assistant in standardized conditions.

3.5 Data Analysis

Repeated measures analysis of variance (ANOVA) will be used to evaluate the effects of the mindful drawing intervention on depressive symptoms at different time points. Main effects of time, group, and their interaction will be analyzed (Field, 2013). For significant results, post hoc tests will be conducted to determine differences between specific time points. The significance level is set at 0.05, and all statistical analyses will be performed using SPSS 26.0.

4. Predicted Results

4.1 Intervention Effects

It is predicted that ANOVA results will reveal significant differences between the intervention and control groups in terms of symptom improvement. It is expected that the intervention group will have significantly lower depressive symptom scores at post-intervention and follow-up (T2 and T3), suggesting that the mindful drawing intervention helps alleviate depressive symptoms.

4.2 Other Possible Findings

In addition to the reduction in depressive symptoms, the study may also find that mindful drawing has positive effects on other psychological indicators, such as anxiety and life satisfaction. It is anticipated that anxiety levels in the intervention group will decrease, and life satisfaction scores will increase. These findings would contribute to a broader understanding of the overall mental health benefits of mindful drawing.

5. Discussion

5.1 Mechanisms

The potential results of this study may indicate that mindful drawing alleviates depressive symptoms through mechanisms such as emotional regulation and self-expression. Mindful drawing may help participants become more aware of and recognize their emotions, thereby improving their ability to regulate negative emotions (Gross, 2015). Additionally, drawing as a form of self-expression may allow participants to externalize their emotions, reduce emotional suppression, and thus promote mental health.

5.2 Limitations and Future Directions

The main limitation of this study is that it remains in the theoretical stage and has not yet undergone practical experimental validation. Therefore, the conclusions drawn from this study require further empirical research to support them. Future research could experimentally validate the effectiveness of mindful drawing and expand the sample size to increase the generalizability of the results. Additionally, future studies could explore the effects of different types of mindfulness-based art interventions on depression and other psychological symptoms to provide more diverse clinical intervention options.

6. Conclusion

This study emphasizes the potential efficacy of mindful drawing in alleviating depressive symptoms. Mindful drawing may help patients reduce depressive emotions and improve mental health through emotional regulation and self-expression. These findings provide a simple yet effective auxiliary intervention for clinical practice, particularly suitable for patients resistant to traditional psychotherapy or those with difficulties in emotional

expression.

Conflict of Interest Statement

The authors declare that there is no conflict of interest regarding the publication of this paper. All research conducted adhered to the highest standards of scientific integrity and ethical practice.

References

1. Baer, R. A. (2003). Mindfulness training as a clinical intervention: a conceptual and empirical review. *Clinical psychology: Science and practice*, 10(2), 125.
2. Campbell, D. T., & Stanley, J. C. (2015). Experimental and quasi-experimental designs for research. Ravenio books.
3. Chambers, R., Gullone, E., & Allen, N. B. (2009). Mindful emotion regulation: An integrative review. *Clinical psychology review*, 29(6), 560-572.
4. Cohen, G. L., & Sherman, D. K. (2014). The psychology of change: Self-affirmation and social psychological intervention. *Annual review of psychology*, 65(1), 333-371.
5. Csikszentmihalyi, M. Flow—the psychology of optimal experience. 1990.
6. Cuijpers, P., Karyotaki, E., Weitz, E., Andersson, G., Hollon, S. D., & van Straten, A. (2014). The effects of psychotherapies for major depression in adults on remission, recovery and improvement: a meta-analysis. *Journal of affective disorders*, 159, 118-126.
7. Ferrari, A. J., Charlson, F. J., Norman, R. E., Patten, S. B., Freedman, G., Murray, C. J., ... & Whiteford, H. A. (2013). Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS medicine*, 10(11), e1001547.
8. Field, A. (2013). Discovering statistics using IBM SPSS statistics.
9. Gross, J. J. (2015). Emotion regulation: Current status and future prospects. *Psychological inquiry*, 26(1), 1-26.
10. Gussak, D. (2007). The effectiveness of art therapy in reducing depression in prison populations. *International Journal of Offender therapy and comparative Criminology*, 51(4), 444-460.
11. Peterson, C., & Rappaport, L. (2014). Mindfulness-based art therapy. Mindfulness and the art therapies: Theory and practice, 42-58.
12. Jakobsen, J. C., Katakam, K. K., Schou, A., Hellmuth, S. G., Stallknecht, S. E., Leth-Møller, K., ... & Gluud, C. (2017). Selective serotonin reuptake inhibitors versus placebo in patients with major depressive disorder. A systematic review with meta-analysis and Trial Sequential Analysis. *BMC psychiatry*, 17, 1-28.
13. Jon Kabat-Zinn, & University of Massachusetts Medical Center/Worcester. Stress Reduction Clinic. (1990). Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness. Delta.
14. Kazdin, A. E., & Tuma, A. H. (1982). Single-case research designs.
15. Keller, M. B., McCullough, J. P., Klein, D. N., Arnow, B.,

- Dunner, D. L., Gelenberg, A. J., ... & Zajecka, J. (2000). A comparison of nefazodone, the cognitive behavioral-analysis system of psychotherapy, and their combination for the treatment of chronic depression. *New England journal of medicine*, 342(20), 1462-1470.
16. Keng, S. L., Smoski, M. J., & Robins, C. J. (2011). Effects of mindfulness on psychological health: A review of empirical studies. *Clinical psychology review*, 31(6), 1041-1056.
17. Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine*, 16(9), 606-613.
18. King, J. L. (2015). Art Therapy: A Brain-based Profession. *The Wiley handbook of art therapy*, 77-89.
19. Monti, D. A., Peterson, C., Kunkel, E. J. S., Hauck, W. W., Pequignot, E., Rhodes, L., & Brainard, G. C. (2006). A randomized, controlled trial of mindfulness-based art therapy (MBAT) for women with cancer. *Psycho-Oncology: Journal of the Psychological, Social and Behavioral Dimensions of Cancer*, 15(5), 363-373.
20. Paykel, E. S. (2009). Residual symptoms and relapse in depression. *Medicographia*, 31(2), 157-163.
21. Segal, Z., Williams, M., & Teasdale, J. (2012). Mindfulness-based cognitive therapy for depression. Guilford press.
22. Steger, M. F., Dik, B. J., & Duffy, R. D. (2012). Measuring meaningful work: The work and meaning inventory (WAMI). *Journal of career Assessment*, 20(3), 322-337.
23. Steger, M. F., Frazier, P., Oishi, S., & Kaler, M. (2006). The meaning in life questionnaire: assessing the presence of and search for meaning in life. *Journal of counseling psychology*, 53(1), 80.
24. World Health Organization. (2017). Depression and other common mental disorders: global health estimates.
25. World Medical Association. (2013). World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *Jama*, 310(20), 2191-2194.