



Current Trends: Recent Innovation and Impacts of Flap Necrosis in Breast Reduction

A. Mohamed Sikkander^{1*}, C. Hazarathaiah Yadav², Narendra Revanuri²

¹Department of Chemistry, Velammal Engineering College, Chennai-600066 INDIA

²Department of Chemistry, Vel Tech Rangarajan Dr. Sakunthala R & D Institute of science & Technology, Avadi, Chennai, India.

*Corresponding Author

A. Mohamed Sikkander

Department of Chemistry,
Velammal Engineering
College, Chennai-600066
INDIA.

Article History

Received: 06.08.2025

Accepted: 10.10.2025

Published: 27.11.2025

Abstract: - Breast obese necrosis is an unsupported fat tissue pain caused by oxygen dissipation made available towards fat cells, ultimately primarily towards cell death. It often occurs in women who endure breast measures, even if they are also able to be present in combination with melanoma. Almost common etiology is disturbing in non-surgical patients. This hustle and bustle review the etiology, pathway physiology, evaluation and relationship of breast fat necrosis surrounded by surgical patients. Pay attention to the permanent times connected by breast fat necrosis. Portray has a characteristic imaging conclusion related to breast fat necrosis. Assess the importance of an alliance and message inside between professional composition members and patients to get results for surgical patients who are intensifying breast fat necrosis. Breast fat necrosis is inanimate fat cells from an offended or is chemic thoracic bandana with the intention of replacing it with a bandana flaw and being present as a glaring prominence.

Keywords: breast fat necrosis, etiologies, melanoma, mammary lymph nodes, fibrous connective tissue.



This is an open access article under the [CC BY-NC](https://creativecommons.org/licenses/by-nc/4.0/) license



Introduction:

Creatures have the two most important types of fat among the characteristic functions: BAT and WAT brown fat tissue [1]. Adult adipocytes in BAT are full of numerous miniature lipid droplets and well above the ground number of mitochondria, while adult adipocytes in WAT are full of small amounts of mitochondria and one-eyed drops of lipids for storing triglycerides [2]. BAT is to produce heat from lipolytic progression with the intention of disentangling the mitochondrial slope of the proton from the ATP cohort [3]. Unlike triglycerides in WAT, they go crazy and ecstatic for tissues that are used as energy throughout oxidation. Characteristically, white adipose tissue (WAT) was well thought out as a stationary component of connective tissue, but is currently valued as the most important monitor of metabolic physiology and hormonal homeostasis [4]. Contemporary works necessary for the development and addition of WAT in vivo highlight the consequences of unequivocal situations of WAT or depositories in the Metabolic Directive [5].

Humorous, adult white adipocytes are incorporated into many tissues. An innovative point of view regarding in vivo guidelines and the WAT target in these tissues has been decked with important responsibility for adipocytes in tissue homeostasis and the Renaissance [6]. Finally, there has been significant progress in the indulgent way in which adult adipocytes confirm the pathology of quite many diseases [7]. In this valuation, we discuss these invented roles of WAT in homeostasis and the renaissance of epithelium, muscles, and immune tissues, and how they enter the pathology of quite many syndromes [8].

Breast fat necrosis has many etiologies and implications; therefore, careful, lasting time is very important to properly assess the patient [9]. Almost all common etiology of fat necrosis is current breast surgery; on the other hand, inoperable patients, melanoma or suffering from a cursory of the thoracic tissue belongs to the perpetrator [10] (**Figure: 1**).

Cite this article:

Sikkander, A. M., Yadav, C. H., Revanuri, N., (2025). Current Trends: Recent Innovation and Impacts of Flap Necrosis in Breast Reduction. *ISAR Journal of Medical and Pharmaceutical Sciences*, 3(11), 12-19.




Figure: 1. Symptom's of Breast Fat Necrosis

Research Methodology:

Pectoral fat necrosis can be successful by intellectual about imaging your breasts with melanoma. It is capable to be cosmetically additionally. In this document, the strength of the form contact point for breast fat necrosis in a surgical patient (Figure: 2).




Figure: 2. Mammography in CC left projection: Multiple calcified lip necrotic post-traumatic cysts & Mammography in right MLO projection: Large calcified lip necrotic trauma cyst.

Three structures include breast: skin, breast tissue and subcutaneous tissue. Breast tissue covers epithelial and stomal elements in cooperation, with fatty and fibrous connective tissue [11] (Figure: 3).




Figure: 3. Schematized observation of the tissue association within breast in the midst of the skin texture describes. Breast tissue overlies the ribs along with torso muscles. The milk

producing glandular epithelia of a Female's breast is enclosed surrounded by adipocyte tissue.

The necessary stromal components are responsible for the advantage of breast capacity in the absence of non-lactation situation [12] (Figure: 4).




Figure: 4. The mammary gland. Schematic representation of adult human mammary gland anatomy. Schematic representation of a terminal end bud (TEB).

The advantage of blood towards the breast appears on or after the perforators of the internal blood vessel of the mammal, in addition to the alternative to lateral thoracic blood vessel perforators [13] (Figure: 5 and 6).




Figure: 5. Illustration of the thoracodorsal vessels arising from the subscapular system




Figure: 6. Disclosure of the thoracodorsal artery, vein, and nerve proceeding to their ingress during in the direction of the latissimusdorsi muscle. The nerve be supposed to be dissect complimentary to put off whichever kinking of the vessels subsequent to anastomosis

This problem is required, especially in the case of breast reduction and / or modernization, somewhere in the vessel transaction and after comparative ischemia be able to show the path to breast fat necrosis [14]. Lymphatic drainage occurs after one-way ejaculation, as in unfathomable subcutaneous and intra-massage vessels, in the direction of the axillaries in addition to the internal lymph nodes [15] (**Figure: 7**).




Figure: 7. Lymphatic drainage of the breast

Although the blood improvement appears in the mammalian internal blood vessel, only 4% of lymph flow goes towards the internal mammary nodes, using a 97% advantage of lymph movements towards axillaries nodes [16](**Figure: 8**).




Figure: 8. Lymph flows in the direction of the axillary's nodes

Result and Discussion:

Fat necrosis can be diagnosed clinically or rasio graphically in the predominance of personal belongings, without the necessity for a biopsy. In surgical patients who have recently undergone a surgical course of breast function, such as breast reduction, modernization includes exclusion or fat transplant after the most important reconstruction, almost everyone has spread their appearance, finding a palpable mass or lump under the skin of the breast [17] (**Figure: 8**).




Figure: 9. Different levels of fat necrosis: i most important level of fat necrosis indicates wreckage of adipose tissue. ii. Primary level of fat necrosis indicates entity adipocytes. iii. Intermediate level of fat necrosis indicates permeation by histiocytes. iv. Intermediate level of fat necrosis indicates conglomeration of RBCs referred to as "mysospherulosis". v. Belatedly stage of fat necrosis indicates single multinucleated giant cell. vi. Belatedly stage of fat necrosis indicates calcifications (are common in late stage of fat necrosis). vii. Belatedly stage of fat necrosis indicates macrophages containing hemosiderin. viii. Belatedly stage of fat necrosis indicates calcifications

There is a preference for subareolar- and periodontal regions, nonetheless it can follow anywhere. Detection by imaging devoid of an observable source, such as current surgery or suffering, or at what time combined by a conclusion, such as lymphadenopathy or skin revolution, necessitate staying away from melanoma [18].

Fat necrosis is almost all often a significance of the breast strain (21 to 70%), very well aspiration or biopsy of the pine needle, emission of anticoagulant performance and breast infection. Patient personality factors such as smoking, corpses, except for the adult era, because the fight is smooth because the action combined with breast melanoma, be particularly combined by increasing breast fat necrosis [19]. Breast fat necrosis is able to embrace the alliance with several breast surgery; on the other hand, it becomes the most important fear of mastectomy / restoration, because fat necrosis is able to cause malformations or breast anxiety anticipated for the re-emergence of cancer [20].

Unladed flutter has gained an act of kindness proposed for the modernization of breasts after mastectomy, exceeding the precedent for several decades [21]. The blood bandana relocation flap contributes to or after a single corpse component to one more. After incomplete or complete mastectomy, this progression is an antilogies breast reconstruction and can also be active due to peduncle or free change of tissue assignment [22].

When moving bandanas of relocate, blood is earmarked in one piece; in addition to fluttering the bandanas is a tunnel or turn to the beneficiary's side. Inside the free excitement, the vessel is

turned off during transfer, resulting in assembly „free“, “and then re-connected micro surgically towards the new-fashion blood vessel except the vein from the side or in close proximity towards the beneficiary's place. Deeply worse epigastria perforator flap (DIEP) is a modern conventional alternative designed for flap without breasts [23].

The establishment of flutter is multi-faceted and necessitates a significant level of teaching, configuration and surgical proficiency [24]. Numerous factors indicate the path towards fat necrosis in the renovation of flutter [25]. On the other hand, the cause of the problem is ischemia - which is a more insufficient main influx or free of venous hemorrhage, is able to justify ischemia in addition to the next fat necrosis with dither alone. Numerous studies have shown the relationship of breast fat necrosis in renovation without any flap costs using the number of perforators and the weight of dither. This conclusion is instinctive because the relative amount of blood delivered to the surface increases, there is additional fat perfusion in the presence and less ischemia [26].

It is also recommended that more venous hemorrhage with added venous anatomists reduce the occurrence of breast fat necrosis. An additional, factor exposed to increased breast fat necrosis in the free flaps are smoke, peduncle caliber, flap type, radiation and the surgeon's occurrence. Many surgeons have need of patients to leave smoking 8 weeks before surgery [27].

There are many techniques presented during breast reduction, based on scratches and peduncle types [28]. Breast reduction is also likely by liposuction and anyone other than intended is mostly kept in reserve for young women with predictable skin elasticity well above ground [29]. Szypulka is a region rich in haberdashery with the help of lonely or numerous main perforators who bought smaller capillaries. The provision of post-postulate blood can be significantly disturbed, especially in remote areas of the cervix in addition to the skin lying above it, after which fat necrosis can occur [30]. The amount of excise duty on breast tissue in a straight line correlates with the obstacle indicator. The occurrence of fat necrosis in breast reduction causes 1 to 9% [31].

Tobacco smoke is a recognized self-determining problem in greater than against the risk of fat necrosis, in addition to the additional complication after breast reduction. The fat graft is supplemented by fat collection by liposuction on or after one body component and injects it into an additional [32]. Fat grafting can be particularly constructive in giving shape malformations in breast reconstruction, known as „gradual deformation“, "or peak anywhere, in which the breast renovation passes towards the resident's chest wall. In addition, a fat transplant is able to achieve better skin quality in the vicinity after emission or mastectomy with a tight fist, reduce visibility in the event of unexplained by means of a small subcutaneous fat or slight resident's chest bandana, in addition to supplementing the cause of the defect by removing previous breast fat necrosis [33].

Fat necrosis is a commonly detected significance after fat transplantation because blood introduces into a fat transplant is accidental and is acquired from neighboring tissues through spread and revascularization. Although numerous fat transplant techniques include an assessment in which it participates, there is no compromise on the dominance in the middle of them. Despite the techniques, the reported fat necrosis pricelist after fat vaccination is surrounded by 2% and 18% [34]. No one can change to see that the

era, body mass index, melanoma stage, smoke, radiation, harvesting site or technological variables come towards predicting the development of fat necrosis [35]. On the other hand, the conducted research shows that the blurring of obesity correlates straight with the time spent ex vivo; therefore, lipo transfer should take place so punctually, as possible after entering fat [36](Figure: 10).




Figure: 10. Fat graft is haphazard and acquire as of adjacent tissues through dissemination and revascularization.

Breast fat necrosis is excellent over and over again because there is a small amount of blood-free adipose at the back, which causes ischemic necrosis. Accumulation is usually expressionless, in addition to the amount of summary of differences, taking into account fat necrosis, myoma, hematoma or seroma, calcification of stitches or skins, scar, swelling, abscess, and also for recurrent or new-fashioned breast cancer. It is necessary to evaluate these patients, because limited re-emergence of melanoma usually occurs in the initial 1 – 5 years after surgery, while most of them have fat necrosis within a few weeks or months after surgery [37,38].

The flaps practice the advanced speed of fat necrosis during irradiation [39,40]. The speediness of indicative fat necrosis in branched therapy in addition to accelerating unfinished breast irradiation is determined towards life from 1 to 50%. Occurrence is associated with dimensions involving a predetermined amount of radiation or how the tissue has been largely irradiated by the highest radiation force previously established [41]. There is a number of verifications in the presence of whether interstitial brachytherapy can provide additional suffering outstanding towards a rooted pine needle, causing greater than before the prevalence of fat necrosis. Studies have also revealed an increase in breast fat necrosis during brachytherapy in populations that also conventionally treat Adriamycin-based chemotherapy [42].

Flap necrosis in mastectomy is a major problem in analgesic reconstruction [43]. Necrosis can perhaps create inconspicuous scars, fabricate contour irregularities and distort the pectoral mound. This may probably show the way to poor modernization and patient satisfaction. Significantly advantage, enlargement and healing of mastectomy flap necrosis may hinder auxiliary oncological healing [44]. I carry out a demonstration assessment of the visual assistance of every patient undergoing antiglare breast modernization within 5 years to examine, occur and cure mastectomy flap necrosis. This information is to produce a feeding algorithm for mastectomy flap necrosis [45,46].

The objectives of this algorithm work are: (a) do not hinder auxiliary oncological treatment, (b) to accelerate therapeutic occurrence at the same time to minimize the risk to the patient, and (c) for aesthetic breast reproduction. The mastectomy housing necrosis is 30% [47]. In the face of large surface area perosis, mastectomy was not delayed in adjuvant oncological treatment [48]. Examination of demonstration records is acceptable to then

build a surveillance algorithm designed for mastectomy housing necrosis. An advantage for the patient and reconstructive is the healing of breast wounds in the acute phase (within 3 weeks); and with regression analysis, we found, that full thickness wounds greater than 6 cm benefit from operational intervention [49]. Finally, patients requiring adjuvant oncological treatment should be cured as soon as possible so that they can carry on oncological treatment [50-64].

Conclusions:

Breast fat necrosis can be determined immediately during postoperative surgery or months after surgery or in difficult breast situations. With the help of hospitalized patients, not only the extravagance of the symptom of persistence is significant, but also the maintenance of the moving saddle in the intelligence with the intention of cep defeat or the cause of breast malformations. Such complications may require a permanent journey towards the business space. In an outpatient environment, it is important for every care provider to become familiar with the risk factor and talk to the patient's care. Comfort, in addition to measurable notes up close, is essential in what is thought out, and proposed for survival by breast fat necrosis.

For the most part, methodical, competent permission is to be part of a pre-operative visit. This conversation would include the wide-ranging risk associated with several operations, other than should also include the likely result and precise complication for breast surgery. Patient memory and appreciation of competent approval can be over and over again deprived of, mainly nervousness and sensations related to breast surgery. For this reason, it is necessary to think about the connection with the competent consent of the conversation about the risk of fat necrosis, hypertrophic development of the spout, wobbly, serious compassion / lack of feeling, aesthetic irregularity, skin necrosis, in addition to / otherwise, as well as devastating the flaps, anyone that may affect the re-visiting of the operating room. In addition, it is necessary to talk about how the strength of the nature of the healthcare unit appeals to these complications; not only is this intended for new complete information, agreement, other than this may further increase the possible permanent satisfaction and knowledge through these results.

Conflict of interest: The authors declare that there are no conflicts of interest

References

1. Townsend, K. L., & Tseng, Y. (2012). Brown adipose tissue. *Adipocyte*, 1(1), 13–24. <https://doi.org/10.4161/adip.18951>
2. Lee, J. H., Park, A., Oh, K., Lee, S. C., Kim, W. K., & Bae, K. (2019). The role of adipose tissue mitochondria: Regulation of mitochondrial function for the treatment of metabolic diseases. *International Journal of Molecular Sciences*, 20(19), 4924. <https://doi.org/10.3390/ijms20194924>
3. Shinde, A. B., Song, A., & Wang, Q. A. (2021). Brown adipose tissue heterogeneity, energy metabolism, and beyond. *Frontiers in Endocrinology*, 12. <https://doi.org/10.3389/fendo.2021.651763>
4. Paul, A., Chanclón, B., Brännmark, C., Wittung-Stafshede, P., Olofsson, C. S., Asterholm, I. W., & Parekh, S. H. (2023). Comparing lipid remodeling of brown adipose tissue, white adipose tissue, and liver after one-week high fat diet intervention with quantitative Raman microscopy. *Journal of Cellular Biochemistry*, 124(3), 382–395. <https://doi.org/10.1002/jcb.30372>
5. Sahu, B., Tikoo, O., Pati, B., Senapati, U., & Bal, N. C. (2022). Role of distinct fat depots in metabolic regulation and pathological implications. In *Reviews of physiology, biochemistry and pharmacology* (pp. 135–176). https://doi.org/10.1007/112_2022_73
6. Machado, S. A., Pasquarelli-Do-Nascimento, G., Da Silva, D. S., Farias, G. R., De Oliveira Santos, I., Baptista, L. B., & Magalhães, K. G. (2022). Browning of the white adipose tissue regulation: new insights into nutritional and metabolic relevance in health and diseases. *Nutrition & Metabolism*, 19(1). <https://doi.org/10.1186/s12986-022-00694-0>
7. Longo, M., Zatterale, F., Naderi, J., Parrillo, L., Formisano, P., Raciti, G. A., Béguinot, F., & Miele, C. (2019). Adipose tissue dysfunction as determinant of Obesity-Associated metabolic complications. *International Journal of Molecular Sciences*, 20(9), 2358. <https://doi.org/10.3390/ijms20092358>
8. Kobayashi, T., Naik, S., & Nagao, K. (2019). Choreographing immunity in the skin epithelial barrier. *Immunity*, 50(3), 552–565. <https://doi.org/10.1016/j.immuni.2019.02.023>
9. Kerridge, W. D., Kryvenko, O. N., Thompson, A., & Shah, B. A. (2015). Fat Necrosis of the Breast: A Pictorial Review of the Mammographic, Ultrasound, CT, and MRI Findings with Histopathologic Correlation. *Radiology Research and Practice*, 2015, 1–8. <https://doi.org/10.1155/2015/613139>
10. Kerridge, W. D., Kryvenko, O. N., Thompson, A., & Shah, B. A. (2015). Fat Necrosis of the Breast: A Pictorial Review of the Mammographic, Ultrasound, CT, and MRI Findings with Histopathologic Correlation. *Radiology Research and Practice*, 2015, 1–8. <https://doi.org/10.1155/2015/613139>
11. Biswas, S. K., Banerjee, S., Baker, G. W., Kuo, C., & Chowdhury, I. (2022). The Mammary Gland: Basic Structure and Molecular Signaling during Development. *International Journal of Molecular Sciences*, 23(7), 3883. <https://doi.org/10.3390/ijms23073883>
12. Sardana, R., Butta, H., Gilotra, K., Mendiratta, L., Baweja, P., & Sarin, R. (2022). Non-diphtheriae corynebacteria causing breast abscess in non-lactating females seeking to avoid pitfalls in diagnosis of chronic breast infections through diagnostic stewardship. *Annals of Pathology and Laboratory Medicine*, 9(6), A115–120. <https://doi.org/10.21276/apalm.3162>
13. Caouette-Laberge, L., & Borsuk, D. E. (2013). Congenital anomalies of the breast. *Seminars in Plastic Surgery*, 27(01), 036–041. <https://doi.org/10.1055/s-0033-1343995>
14. Kaidar-Person, O., Eblan, M. J., Caster, J. M., Shah, A., Fried, D., Marks, L. B., Lee, C. N., & Jones, E. L. (2019). Effect of internal mammary vessels radiation dose on outcomes of free flap breast reconstruction. *the αBreast Journal*, 25(2), 286–289. <https://doi.org/10.1111/tbj.13202>
15. Sardana, R., Butta, H., Gilotra, K., Mendiratta, L., Baweja, P., & Sarin, R. (2022b). Non-diphtheriae corynebacteria causing

- breast abscess in non-lactating females seeking to avoid pitfalls in diagnosis of chronic breast infections through diagnostic stewardship. *Annals of Pathology and Laboratory Medicine*, 9(6), A115-120. <https://doi.org/10.21276/apalm.3162>
16. Abstracts critique - IJCCM2021. (2021). *Indian Journal of Critical Care Medicine/Indian Journal of Critical Care Medicine*, 25(S1), S1–S114. <https://doi.org/10.5005/ijccm-25-s1-s1>
17. Professional, C. C. M. (n.d.). Fat necrosis. Cleveland Clinic. <https://my.clevelandclinic.org/health/diseases/24187-fat-necrosis>
18. Bayram, Y., Sezgiç, M., Karakol, P., Bozkurt, M., & Fılinte, G. T. (2019). The use of autologous fat grafts in breast surgery: A literature review. *Archives of Plastic Surgery*, 46(06), 498–510. <https://doi.org/10.5999/aps.2019.00416>
19. Demark-Wahnefried, W., Peterson, B. L., Winer, E. P., Marks, L. B., Aziz, N. M., Marcom, P. K., Blackwell, K., & Rimer, B. K. (2001). Changes in weight, body composition, and factors influencing energy balance among premenopausal breast cancer patients receiving adjuvant chemotherapy. *Journal of Clinical Oncology*, 19(9), 2381–2389. <https://doi.org/10.1200/jco.2001.19.9.2381>
20. Lee, J., Park, H. Y., Kim, W. W., Lee, J. J., Keum, H. J., Yang, J. D., Lee, J. W., Lee, J. S., & Jung, J. H. (2021). Natural course of fat necrosis after breast reconstruction: a 10-year follow-up study. *BMC Cancer*, 21(1). <https://doi.org/10.1186/s12885-021-07881-x>
21. Agarwal, S., Pappas, L. M., Neumayer, L., Kokeny, K. E., & Agarwal, J. (2014). Effect of Breast Conservation Therapy vs Mastectomy on Disease-Specific Survival for Early-Stage Breast Cancer. *JAMA Surgery*, 149(3), 267. <https://doi.org/10.1001/jamasurg.2013.3049>
22. Garza, R., Ochoa, O., & Chrysopoulou, M. (2021). Post-mastectomy Breast Reconstruction with Autologous Tissue: Current Methods and Techniques. *Plastic and Reconstructive Surgery. Global Open*, 9(2), e3433. <https://doi.org/10.1097/gox.0000000000003433>
23. Allen, R. J., & Treece, P. (1994). Deep inferior epigastric perforator flap for breast reconstruction. *Annals of Plastic Surgery*, 32(1), 32–38. <https://doi.org/10.1097/0000637-199401000-00007>
24. Kahol, K., Satava, R. M., Ferrara, J. J., & Smith, M. (2009). Effect of Short-Term pretrial practice on surgical proficiency in simulated environments: A randomized trial of the “Preoperative Warm-Up” effect. *Journal of the American College of Surgeons*, 208(2), 255–268. <https://doi.org/10.1016/j.jamcollsurg.2008.09.029>
25. Fat necrosis. (n.d.). <https://www.pathologyoutlines.com/topic/breastfatnecrosis.html>.
26. Mulvey, C. L., Cooney, C. M., Daily, F. F., Colantuoni, E., Ogbuago, O. U., Cooney, D. S., Rad, A. N., Manahan, M. A., Rosson, G. D., & Sacks, J. M. (2013). Increased flap weight and decreased perforator number predict fat necrosis in DIEP breast reconstruction. *Plastic and Reconstructive Surgery. Global Open*, 1(2), 1–7. <https://doi.org/10.1097/gox.0b013e318294e41d>
27. Metz, G., Snook, K., Sood, S., Baron-Hay, S., Spillane, A., Lamoury, G., & Carroll, S. (2022). Breast Radiotherapy after Oncoplastic Surgery—A Multidisciplinary Approach. *Cancers*, 14(7), 1685. <https://doi.org/10.3390/cancers14071685>
28. Grover, K., Marano, A. A., Lin, A. J., Peysakhovich, A., Castillo, W., & Rohde, C. H. (2023). The modified Wise-pattern superomedial pedicle reduction mammoplasty for benign macromastia: a 9-year case series. *Annals of Breast Surgery*, 7, 14. <https://doi.org/10.21037/abs-21-102>
29. Moskovitz, M. J., & Baxt, S. A. (2004). Breast reduction using liposuction alone. *Seminars in Plastic Surgery*, 18(3), 225–229. <https://doi.org/10.1055/s-2004-831909>
30. Yoon, C. S., & Kim, K. N. (2016). Selective salvage of zones 2 and 4 in the pedicled TRAM flap: a focus on reducing fat necrosis and improving aesthetic outcomes. *SpringerPlus*, 5(1). <https://doi.org/10.1186/s40064-016-1714-7>
31. Klimberg, V. S., & Rivere, A. (2016). Ultrasound image-guided core biopsy of the breast. *Chinese Clinical Oncology*, 5(3), 33. <https://doi.org/10.21037/cco.2016.04.05>
32. The MNT Editorial Team. (2018, November 16). What is medicine? <https://www.medicalnewstoday.com/articles/323679#1>
33. Rohrich, R. J., Ghavami, A., Constantine, F. C., Unger, J. G., & Mojallal, A. (2014). Lift-and-Fill face lift. *Plastic and Reconstructive Surgery/PSEF CD Journals*, 133(6), 756e–767e. <https://doi.org/10.1097/01.prs.0000436817.96214.7e>
34. Turner, A., Abu-Ghname, A., Davis, M. J., Winocour, S., Hanson, S. E., & Chu, C. K. (2020). Fat grafting in breast reconstruction. *Seminars in Plastic Surgery*, 34(01), 017–023. <https://doi.org/10.1055/s-0039-1700959>
35. Noszczyk, B., & Krześniak, N. (2013). Fat grafts in the reconstruction and treatment of chronic wounds. *Polski Przegląd Chirurgiczny*, 85(12). <https://doi.org/10.2478/pjs-2013-0113>
36. The 10th Asia-Pacific Primary Liver Cancer Expert Meeting (APPLE 2019), Sapporo, Japan, August 29-31, 2019. (2019). *Liver Cancer*, 8(Suppl. 1), 1–204. <https://doi.org/10.1159/000502497>
37. Kølle, S. T., Duscher, D., Taudorf, M., Fischer-Nielsen, A., Svalgaard, J. D., Munthe-Fog, L., Jønsson, B., Selvig, P. B., Mamsen, F. P. W., & Katz, A. J. (2020). Ex vivo-expanded autologous adipose tissue-derived stromal cells ensure enhanced fat graft retention in breast augmentation: A randomized controlled clinical trial. *Stem Cells Translational Medicine*, 9(11), 1277–1286. <https://doi.org/10.1002/sctm.20-0081>
38. Brambullo, T., Azzena, G. P., Toninello, P., Masciopinto, G., De Lazzari, A., Biffoli, B., Vindigni, V., & Bassetto, F. (2021). Current surgical therapy of locally advanced CSCC: From patient selection to microsurgical tissue transplant.

- review. *Frontiers in Oncology*, 11. <https://doi.org/10.3389/fonc.2021.783257>
39. Tol, J. a. M. G., Gouma, D. J., Bassi, C., Dervenis, C., Montorsi, M., Adham, M., Andrén-Sandberg, Å., Asbun, H. J., Bockhorn, M., Büchler, M. W., Conlon, K. C., Fernández-Cruz, L., Fingerhut, A., Friess, H., Hartwig, W., Izbicki, J. R., Lillemoe, K. D., Milićević, M., Neoptolemos, J. P., . . . Charnley, R. (2014). Definition of a standard lymphadenectomy in surgery for pancreatic ductal adenocarcinoma: A consensus statement by the International Study Group on Pancreatic Surgery (ISGPS). *Surgery*, 156(3), 591–600. <https://doi.org/10.1016/j.surg.2014.06.016>
40. Kerridge, W. D., Kryvenko, O. N., Thompson, A., & Shah, B. A. (2015c). Fat Necrosis of the Breast: A Pictorial Review of the Mammographic, Ultrasound, CT, and MRI Findings with Histopathologic Correlation. *Radiology Research and Practice*, 2015, 1–8. <https://doi.org/10.1155/2015/613139>
41. Clarke-Pearson, E. M., Chadha, M., Dayan, E., Dayan, J. H., Samson, W., Sultan, M. R., & Smith, M. L. (2013). Comparison of Irradiated Versus Nonirradiated DIEP Flaps in Patients Undergoing Immediate Bilateral DIEP Reconstruction with Unilateral Postmastectomy Radiation Therapy (PMRT). *Annals of Plastic Surgery*, 71(3), 250–254. <https://doi.org/10.1097/sap.0b013e31828986ec>
42. Lee, J., Park, H. Y., Kim, W. W., Lee, J. J., Keum, H. J., Yang, J. D., Lee, J. W., Lee, J. S., & Jung, J. H. (2021b). Natural course of fat necrosis after breast reconstruction: a 10-year follow-up study. *BMC Cancer*, 21(1). <https://doi.org/10.1186/s12885-021-07881-x>
43. Pagliara, D., Schiavone, L., Garganese, G., Bove, S., Montella, R. A., Costantini, M., Rinaldi, P., Bottosso, S., Grieco, F., Rubino, C., Salgarello, M., & Ribuffo, D. (2023). Predicting Mastectomy skin flap necrosis: A Systematic review of preoperative and intraoperative assessment techniques. *Clinical Breast Cancer*, 23(3), 249–254. <https://doi.org/10.1016/j.clbc.2022.12.021>
44. Matsen, C., Mehrara, B. J., Eaton, A., Capko, D., Berg, A. L., Stempel, M., Van Zee, K. J., Pusic, A. L., King, T. A., Cody, H. S., Pilewskie, M., Cordeiro, P. G., Sclafani, L. M., Plitas, G., Gemignani, M. L., Disa, J. J., El-Tamer, M., & Morrow, M. (2015). Skin flap necrosis after mastectomy with reconstruction: a prospective study. *Annals of Surgical Oncology*, 23(1), 257–264. <https://doi.org/10.1245/s10434-015-4709-7>
45. Luze, H., Nischwitz, S. P., Wurzer, P., Winter, R., Spendel, S., Kamolz, L., & Bjelic-Radisic, V. (2022). Assessment of Mastectomy Skin Flaps for Immediate Reconstruction with Implants via Thermal Imaging—A Suitable, Personalized Approach? *Journal of Personalized Medicine*, 12(5), 740. <https://doi.org/10.3390/jpm12050740>
46. Frey, J. D., Salibian, A. A., Choi, M., & Karp, N. S. (2019). The importance of tissue perfusion in reconstructive breast surgery. *Plastic and Reconstructive Surgery/PSEF CD Journals*, 144(1S), 21S-29S. <https://doi.org/10.1097/prs.0000000000005947>
47. Masood, S. (2016). Neoadjuvant chemotherapy in breast cancers. *Women's Health*, 12(5), 480–491. <https://doi.org/10.1177/1745505716677139>
48. Liu, E. H., Zhu, S., Hu, J., Wong, N., Farrokhyar, F., & Thoma, A. (2019). Intraoperative SPY Reduces Post-mastectomy skin flap Complications: A Systematic Review and Meta-Analysis. *Plastic and Reconstructive Surgery. Global Open*, 7(4), e2060. <https://doi.org/10.1097/gox.0000000000002060>
49. Norman, G., Shi, C., Goh, E. L., Murphy, E. M., Reid, A. J., Chiverton, L., Stankiewicz, M., & Dumville, J. C. (2022). Negative pressure wound therapy for surgical wounds healing by primary closure. *Cochrane Library*, 2022(4). <https://doi.org/10.1002/14651858.cd009261.pub7>
50. Harbeck, N., Penault-Llorca, F., Cortés, J., Gnant, M., Houssami, N., Poortmans, P., Ruddy, K. J., Tsang, J., & Cardoso, F. (2019). Breast cancer. *Nature Reviews. Disease Primers*, 5(1). <https://doi.org/10.1038/s41572-019-0111-2>
51. Deptuła, M., Zieliński, J., Wardowska, A., & Piśkuła, M. (2019). Wound healing complications in oncological patients: perspectives for cellular therapy. *Postępy Dermatologii i Alergologii*, 36(2), 139–146. <https://doi.org/10.5114/ada.2018.72585>
52. Schroen, A. T., & Cress, R. D. (2001). Use of surgical procedures and adjuvant therapy in rectal cancer treatment. *Annals of Surgery*, 234(5), 641–651. <https://doi.org/10.1097/0000658-200111000-00009>
53. Murakami-Nishimagi, Y., Sugimoto, K., Kobayashi, M., Tachibana, K., Kojima, M., Okano, M., Hashimoto, Y., Saji, S., Ohtake, T., & Chiba, H. (2023). Claudin-4-adhesion signaling drives breast cancer metabolism and progression via liver X receptor β . *Breast Cancer Research*, 25(1). <https://doi.org/10.1186/s13058-023-01646-z>
54. Tawfik, E. H., Ghallab, E., & Moustafa, A. a. a. E. (2023). A nurse versus a chatbot – the effect of an empowerment program on chemotherapy-related side effects and the self-care behaviors of women living with breast Cancer: a randomized controlled trial. *BMC Nursing*, 22(1). <https://doi.org/10.1186/s12912-023-01243-7>
55. Vogens, M., Naghavi-Behzad, M., Harbo, F. S. G., Jakobsen, N. M., Gerke, O., Asmussen, J. T., Nissen, H. J., Dahlsgaard-Wallenius, S. E., Braad, P., Jensen, J. D., Ewertz, M., & Hildebrandt, M. G. (2023). 2-[18F]FDG-PET/CT is a better predictor of survival than conventional CT: a prospective study of response monitoring in metastatic breast cancer. *Scientific Reports*, 13(1). <https://doi.org/10.1038/s41598-023-32727-w>
56. Cardoso, F., Paluch-Shimon, S., Senkus, E., Curigliano, G., Aapro, M., André, F., Barrios, C. H., Bergh, J., Bhattacharyya, G., Biganzoli, L., Boyle, F., Cardoso, M., Carey, L. A., Cortés, J., Saghir, N. S. E., Elzayat, M., Eniu, A., Fallowfield, L., Francis, P. A., . . . Winer, E. P. (2020). 5th ESO-ESMO international consensus guidelines for advanced breast cancer (ABC 5). *Annals of Oncology*, 31(12), 1623–1649. <https://doi.org/10.1016/j.annonc.2020.09.010>

57. Sikkander, A. M. (2022). Duct cancer evaluation in situ–review. In *Acta Biology Forum* (pp. 01-04).
58. Sikkander, M., Vedhi, C., & Manisankar, P. (2012). Cyclic voltammetric determination of 1, 4-Dihydro pyridine drugs using MWCNTs modified glassy carbon electrode. *Der Chem. Sin*, 3, 413-420.
59. A. Mohamed Sikkander (2022). Intrathecal Chemotherapy for Blood Cancer Treatment. *Acta Biology Forum*.V01i01, 14-17. DOI: <https://doi.org/10.5281/zenodo.7008901>
60. Sikkander, M., & Nasri, N. S. (2013). Review on Inorganic Nano crystals unique benchmark of Nanotechnology. *Moroccan Journal of Chemistry*, 1(2), J-Chem.
61. Yadav, C. H., Revanuri, N., & Sikkander, A. R. M. (2025). Tungsten-based compounds: A new frontier in cancer diagnosis and therapy. *Journal of Applied Organometallic Chemistry*, 5, 149-167.
62. Sikkander, A. R. M. (2025). RUTHENIUM ORGANOMETALLIC COMPOUNDS IN CANCER TREATMENT. *Biomedical Engineering: Applications, Basis and Communications*, 37(01), 2430003.
63. Yadav,H.C., Revanuri,N., Sikkander,A.R.M, 2025. Organometallic Compounds phototoxicity against cancer cells. *Biomedical Engineering Applications Basis and Communications*, 2550020. <http://dx.doi.org/10.4015/S1016237225500206>
64. Sikkander, A. R. M., Meena, M., Yadav, H., Wahi, N., & Lakshmi, V. V. (2024). Appraisal of the impact of applying organometallic compounds in cancer therapy. *Journal of Applied Organometallic Chemistry*, 4, 145-166.